

Cosmic censorship inside black holes

Lárus Thorlacius*

Science Institute, University of Iceland,

Dunhaga 3, 107 Reykjavk, Iceland

Abstract

A simple argument is given that a traversable Cauchy horizon inside a black hole is incompatible with unitary black hole evolution. The argument assumes the validity of black hole complementarity and applies to a generic black hole carrying angular momentum and/or charge. In the second part of the paper we review recent work on the semiclassical geometry of two-dimensional charged black holes.

*Electronic address: lth@hi.is




FIG. 1: Penrose diagram for maximally extended Reissner-Nordström spacetime. Thick lines represent timelike singularities, dashed lines are event horizons, and dotted lines are Cauchy horizons. The diagram is periodic in the vertical direction with an infinite number of asymptotic regions.

I. INTRODUCTION

A stationary rotating black hole in classical general relativity is described by the maximally extended Kerr spacetime, which has multiple asymptotic regions and black hole regions that contain timelike singularities. For a massive black hole with large angular momentum there exist timelike paths connecting different asymptotic regions without ever entering regions of strong curvature and one can speculate whether macroscopic observers could travel along such paths through the black hole to another “universe”. There are also paths in the Kerr geometry that lead past a singularity into an exotic region with closed timelike curves. The spherically symmetric Reissner-Nordström spacetime of a static electrically charged black hole also has multiple asymptotic regions and timelike singularities but no closed timelike curves. The Penrose diagram of maximally extended Reissner-Nordström spacetime is shown in Figure 1 and, for comparison, the Penrose diagram for an uncharged Schwarzschild black hole is shown in Figure 2.




FIG. 2: Penrose diagram for maximally extended Schwarzschild spacetime. Thick lines represent spacelike singularities and dashed lines are event horizons.

In both the Kerr and Reissner-Nordström cases the physical relevance of much of the extended spacetime has, however, been called into question [1, 2, 3, 4, 5, 6]. Inside each black hole region there is a Cauchy horizon, beyond which a timelike singularity becomes visible. The Cauchy horizon is a surface of infinite blueshift with respect to the event horizon of the black hole in question and this leads to a dynamical instability, referred to as mass inflation, which replaces the Cauchy horizon by a null singularity that turns spacelike deep inside the black hole [7, 8, 9, 10, 11]. However, the null singularity is weak in the sense that integrated tidal effects acting on extended observers are finite and this leaves open the possibility of extending the physical spacetime through it [8]. Such an extension requires ingoing flux of negative energy along the null singularity itself and while negative energy is pathological at the classical level this not necessarily the case in the semiclassical context of quantum field theory in a curved background spacetime.

In view of this it is natural to ask how quantum effects modify the classical geometry inside a rotating or charged black hole. Do they render the Cauchy horizon traversable by supplying the flux of negative energy that is required for extending the geometry? Or, is the Cauchy horizon further destabilized by quantum effects and replaced by a spacelike singularity [4, 12]? In the following we present two very different approaches to this problem that both support the latter possibility. The first one, discussed in Section II, is based on Hawking's black hole information paradox [13]. It is qualitative and rather speculative but applies to generic black holes, with or without angular momentum and charge. The other approach is quantitative but more restricted in scope. It involves charged black holes in a two-dimensional toy model of gravity where quantum effects can be studied systematically [14, 15]. Section III contains a brief review of recent work on this model.

II. A PERSPECTIVE FROM BLACK HOLE COMPLEMENTARITY

In this section we argue that a traversable Cauchy horizon inside a black hole formed in gravitational collapse is incompatible with quantum mechanical unitarity. The argument assumes the validity of the principle of black hole complementarity [16] but is otherwise quite general and applies to both rotating and charged black holes.

A. Unitary black hole evolution

Let us adopt the point of view that the formation and subsequent evaporation of a rotating and/or charged black hole is a unitary process [17, 18]. A number of theoretical developments, including matrix theory [19], the gauge theory/gravity correspondence [20], and the microscopic computation of black hole entropy in string theory [21], support this viewpoint even if the detailed implementation of unitarity has yet to be understood (see [22, 23] for recent discussions of unitarity and the black hole information paradox).

Let us further assume that unitarity is maintained by encoding the information about the initial quantum state of the matter that forms the black hole into subtle correlations in the outgoing Hawking radiation. In principle, this information eventually becomes available to observers outside the black hole although quite hard to come by in practise. The late time Hawking radiation will also contain information about the state of an observer that falls into the black hole during its lifetime but this appears to contradict the expectation, based on the equivalence principle, that an observer in free fall encounters nothing unusual upon crossing the event horizon of a large black hole.

Quantum states cannot be cloned [24, 25] and therefore, as far as distant observers are concerned, any information that emerges in the outgoing Hawking radiation must be removed from the infalling matter before it enters the black hole. On the other hand, from the point of view of an infalling observer who passes unharmed through the event horizon no information has been removed and its duplication in the Hawking radiation appears to violate the principles of quantum mechanics.




FIG. 3: Gedanken experiment for a black hole with a spacelike singularity (thick line). The dashed line is the event horizon and the dotted line represents a null signal from $\mathcal{O}_1$ which is to be received by $\mathcal{O}_2$ before $\mathcal{O}_2$ hits the singularity.

B. Gedanken experiment involving correlated spins

The principle of black hole complementarity states that both viewpoints are equally valid and that apparent contradiction only comes about when we attempt to compare the physical description in the very different reference frames of these observers. In [26] this was illustrated by analysing various gedanken experiments involving black holes designed to expose potential contradictions.

Let us consider one of these experiments which involves quantum correlations between degrees of freedom inside and outside the event horizon. A pair of spins is prepared in a singlet state outside a large black hole and then one of the spins is carried into the black hole by observer $\mathcal{O}_1$ in Figure 3 while the other spin remains outside the black hole and is never actually measured in the experiment. Here “spin” is taken to mean some internal label because conventional spin can in principle be detected at long-range by its gravitational field.

Upon crossing the event horizon, which is indicated by a dashed line in the Figure 3, the infalling observer $\mathcal{O}_1$ carries out a measurement of the spin and transmits the result via a null signal, shown as a dotted line in the figure. Meanwhile observer $\mathcal{O}_2$ makes measurements on the Hawking radiation coming from the black hole. Assuming all information about the quantum state inside the black hole is encoded in the Hawking radiation $\mathcal{O}_2$ can effectively




FIG. 4: Gedanken experiment for a black hole with a traversible Cauchy horizon. In this case $\mathcal{O}_2$ can receive signals after passing through the Cauchy horizon.

measure a component of the spin that went inside the black hole. $\mathcal{O}_2$ then passes inside the event horizon where he can receive the signal from $\mathcal{O}_1$, which potentially contradicts the measurement of the Hawking radiation, in violation of the laws of quantum mechanics.

If the black hole singularity is spacelike, as in Figure 3, there is no paradox here. The information in the Hawking radiation is not immediately available. In fact, less than one bit of information is accessible until the black hole area has evaporated to half its initial value and after that it comes out gradually [27]. Observer $\mathcal{O}_2$ must therefore wait outside the evaporating black hole for a time, which is of order the black hole lifetime, before the measurements of the Hawking radiation can yield any useful information. By then the extreme redshift of the region close to the event horizon, prevents $\mathcal{O}_2$ from learning of the potential contradiction before hitting the singularity. The proper time available for $\mathcal{O}_1$ to carry out the spin measurement and transmit the signal, so that it can be received by $\mathcal{O}_2$, turns out to be much shorter than a Planck time [26].

If, on the other hand, the black hole has a traversable Cauchy horizon, as in Figure 4, then $\mathcal{O}_2$ is at leisure to receive signals after passing through the Cauchy horizon and there is no stringent time limit for $\mathcal{O}_1$ to carry out the spin measurement and transmit the result. A black hole with a traversable Cauchy horizon is therefore incompatible with black hole

complementarity and we conclude that black holes cannot have traversable Cauchy horizons.

We note that the same gedanken experiment was considered in [28] in a Reissner-Nordström background geometry. Those authors did not question the validity of the classical static geometry but instead concluded that the experiment presents a problem for unitarity and the principle of black hole complementarity.

III. TWO-DIMENSIONAL CHARGED BLACK HOLES

In this section we review our recent work with A. Frolov and K. Kristjansson [14, 15] where we study the semiclassical geometry of electrically charged black holes in the simplified context of 1+1-dimensional dilaton gravity coupled to an abelian gauge field. The model has classical charged black hole solutions for which the maximally extended spacetime has multiple asymptotic regions separated by black hole regions containing timelike singularities and associated Cauchy horizons. The Penrose diagram is in fact identical to that of a 3+1-dimensional Reissner-Nordström black hole, shown in Figure 1. Furthermore, mass inflation has been shown to occur at the classical level in 1+1-dimensional dilaton gravity models of this type [29, 30, 31], suggesting that this is indeed a suitable toy model to study the fate of Cauchy horizons at the quantum level.

The main new feature of [14, 15] is to include quantum effects due to electrically charged matter, which turns out to significantly modify the internal geometry of charged black holes in the model. The timelike singularities and Cauchy horizons of a static classical black hole are replaced by a spacelike singularity at the semiclassical level and the global topology becomes the same as that of an electrically neutral Schwarzschild black hole. This conclusion is reached by a combination of analytic and numerical calculations that we only outline here, referring to [14, 15] for a more detailed description.

A. Classical theory

Our starting point is the classical action,

$$S_0 = \int d^2x \sqrt{-g} e^{-2\phi} \left[R + 4(\nabla\phi)^2 + 4\lambda^2 - \frac{1}{4}F^2 \right], \quad (1)$$

with a dilaton field ϕ , an abelian gauge field $F_{\mu\nu}$ and the 1+1-dimensional metric $g_{\mu\nu}$. Due to the factor of $e^{-2\phi}$ in front, the strength of both the gravitational and gauge couplings is

determined by the value of the dilaton field.

The model can be viewed as the s-wave reduction of 3+1-dimensional dilaton gravity in an extremal black hole background [32, 33, 34]. The mass scale λ is proportional to the inverse of the magnetic charge carried by the 3+1-dimensional extremal dilaton black hole. We will use units where $\lambda = 1$. We also note that the two-dimensional dilaton field ϕ has an interpretation in terms of an area in the higher dimensional theory. The precise relation is that the area of the transverse two-sphere in the Einstein frame of 3+1-dimensional dilaton gravity is given by $\psi \equiv e^{-2\phi}$ and thus we refer to ψ as the *area function*.

The action (1) has classical solutions, which are analogous to Reissner-Nordström black holes,

$$\phi = -x, \tag{2}$$

$$ds^2 = -a(x)dt^2 + \frac{1}{a(x)}dx^2, \tag{3}$$

$$F_{tx} = Qe^{-2x}, \tag{4}$$

with

$$a(x) = 1 - Me^{-2x} + \frac{1}{8}Q^2e^{-4x}. \tag{5}$$

The constants M and Q are the mass and charge and for $M > |Q|/\sqrt{2}$ we have a classical black hole geometry with horizons located at the two zeroes of the metric $a(x)$, where the area function takes the following values,

$$\psi_{\pm} = \frac{1}{2} \left(M \pm \sqrt{M^2 - \frac{1}{2}Q^2} \right). \tag{6}$$

The coupling strength e^{ϕ} goes to zero in the asymptotic region $x \rightarrow \infty$ and the metric approaches the 1+1-dimensional Minkowski metric there. On the other hand, there is a curvature singularity inside the black hole, where the area function goes to zero and the coupling diverges.

B. Charged matter

We now add matter to the model to be able to consider dynamical solutions involving gravitational collapse and also to study semiclassical corrections to the geometry due to matter quantum effects. In order to form charged black holes by gravitational collapse we

need to have some form of electrically charged matter and we take this to be charged Dirac fermions,

$$S_m = \int d^2x \sqrt{-g} [i\bar{\Psi}\gamma^\mu(\mathcal{D}_\mu + ieA_\mu)\Psi - m\bar{\Psi}\Psi] \quad (7)$$

where e and m are the fermion charge and mass respectively, and A_μ is the 1+1-dimensional gauge potential.

At the quantum level, fermions will be pair-produced in the electric field of a charged black hole via the Schwinger effect [35] and we are particularly interested in the resulting electromagnetic screening of the black hole charge and the gravitational backreaction on the black hole geometry due to the charged pairs. We are also interested in quantum effects due to electrically neutral matter, *i.e.* Hawking emission [36] of neutral particles from our charged black holes, but let us first consider the effect of Schwinger pair-production and come back to include the Hawking effect later on.

In our two-dimensional model pair-production of charged particles is most conveniently described by bosonizing the fermions. This has two important advantages. First of all, the matter equations of motion are then converted into scalar field equations which are simpler to analyze both analytically and numerically than the original fermion equations. The second advantage is that, since bosonization is a strong-to-weak coupling duality, the quantum mechanical process of fermion pair-production is well described by classical equations for the bosons. This holds when $m \ll e$ and that happens to be the range of parameters of most interest to us because of the very large charge-to-mass ratio of electrons in the real world.

The identification [37, 38] between composite operators of the fermion field and of a real boson field Z carries over from flat to curved spacetime [39], with appropriate replacement of derivatives by covariant derivatives, as long as the background geometry is slowly varying on the microscopic length scale of the matter system. The bosonic description thus breaks down near singularities where the spacetime curvature gets large on microscopic length scales, but this is not a problem for the bosonization *per se* since all classical or semiclassical equations are inadequate in such regions.

The electromagnetic current is given by

$$j^\mu = e\bar{\Psi}\gamma^\mu\Psi = \frac{e}{\sqrt{\pi}}\varepsilon^{\mu\nu}\nabla_\nu Z, \quad (8)$$

where $\varepsilon^{\mu\nu}$ is an antisymmetric tensor related to the Levi-Civita tensor density by $\varepsilon^{\mu\nu} =$

$\epsilon^{\mu\nu}/\sqrt{-g}$. The boson effective action is

$$S_b = \int d^2x \sqrt{-g} \left[-\frac{1}{2}(\nabla Z)^2 - V(Z) - \frac{e}{\sqrt{4\pi}} \epsilon^{\mu\nu} F_{\mu\nu} Z \right], \quad (9)$$

where $V(Z) = c e m (1 - \cos(\sqrt{4\pi} Z))$ and c is an $O(1)$ numerical constant whose precise value will not be needed.

C. Semiclassical black holes

The semiclassical geometry of charged black holes is obtained by solving the equations of motion of the combined dilaton gravity and boson action (1) and (9). We work in conformal gauge $ds^2 = -e^{2\rho} d\sigma^+ d\sigma^-$ and take advantage of the fact that a two-dimensional gauge field is non-dynamical. The two-dimensional Maxwell equations can be used to express the field strength in terms of the dilaton and the bosonized matter field,

$$F^{\mu\nu} = - \left(-\frac{e}{\sqrt{\pi}} Z + q \right) e^{2\phi} \epsilon^{\mu\nu}, \quad (10)$$

where q is a constant of integration which can be interpreted as a background charge located at the strong-coupling end of the one-dimensional space. We are primarily interested in gravitational collapse of charged matter into an initial vacuum and then it is natural to set $q = 0$. We also note that the value of $Z/\sqrt{\pi}$ at a given location determines the amount of electric charge to the left of, or “inside”, that location. This can also be inferred from the bosonized current in (8). It follows that electromagnetic screening will appear as spatial variation in Z in the bosonized formalism.

The remaining semiclassical equations are simplified if we introduce $\theta = 2(\rho - \phi)$ and work with the area function ψ rather than the dilaton itself. This leads to a system of equations,

$$-4\partial_+ \partial_- \psi = \left(4 - \frac{e^2 Z^2}{2\pi\psi^2} \right) e^\theta - \frac{V(Z)e^\theta}{\psi}, \quad (11)$$

$$-4\partial_+ \partial_- \theta = \frac{e^2 Z^2 e^\theta}{\pi\psi^3} + \frac{V(Z)e^\theta}{\psi^2}, \quad (12)$$

$$-4\partial_+ \partial_- Z = \frac{e^2 Z e^\theta}{\pi\psi^2} + \frac{V'(Z)e^\theta}{\psi}, \quad (13)$$

along with two constraints

$$\partial_\pm^2 \psi - \partial_\pm \theta \partial_\pm \psi = -\frac{1}{2}(\partial_\pm Z)^2. \quad (14)$$




FIG. 5: Numerical black hole formation. The figure shows a density plot of the area function ψ for an incoming pulse of charged matter. The 1+1-dimensional spacetime is represented in (compactified) null coordinates. The curvature singularity is indicated by the thick black curve and the apparent horizon by the thin curve. The singularity is spacelike.

In [14, 15] this system was solved numerically for initial data consisting of a left-moving kink configuration in the bosonized matter field, corresponding to incoming charged matter undergoing gravitational collapse. The height of the kink determines the total incoming charge while its steepness governs the total incoming energy.

Figures 5 and 6 show density plots for the area function ψ and the bosonized matter field Z for a typical solution. The geometry is represented in null coordinates and we have performed a conformal compactification in order to cover the entire spacetime in the plots. Curvature singularities, which occur at $\psi = 0$, are indicated by thick black curves and apparent horizons, where $\partial_+ \psi = 0$ or $\partial_- \psi = 0$ [40], by thin black curves.

It is clear from these plots that a charged black hole forms with a curvature singularity inside an apparent horizon. The bosonic matter field goes to zero deep inside the black hole. This reflects charge screening due to fermion pair-production. In fact, for this particular solution the screening effect extends well outside the black hole region. We also note that after the black hole has formed the value of Z decreases along the apparent horizon, *i.e.* the black hole gradually discharges and the final state is a neutral black hole.

The black hole singularity is everywhere spacelike and approaches the apparent horizon




FIG. 6: Numerical evolution of an incoming pulse of charged matter. The density plot shows the bosonized matter field Z , which represents the charge distribution. The white region on the lower left is the vacuum before the arrival of the matter pulse. Charge screening that prevents the Z field from penetrating into the strong-coupling region near the singularity is evident.

at future null infinity. There is no indication of a Cauchy horizon or a null singularity inside the semiclassical black hole.

D. Static solutions

One can also study static black holes in this semiclassical theory. For this we require the fields to only depend on a spatial variable $\sigma = \frac{1}{2}(\sigma^+ - \sigma^-)$ and make a further change of variables, writing $\xi = e^\theta$ and defining a new spatial coordinate via $dy = \xi d\sigma$. In these variables the classical charged black hole solution (2)-(5) can be extended inside the black hole horizon in a straightforward way and the corresponding extension for a semiclassical black hole can be carried out numerically, as described in detail in [14, 15]. The result of such a numerical integration is shown in Figure 7a, with the corresponding classical solution shown in Figure 7b for comparison.

The classical solution has two horizons where $\xi(y)$ goes to zero. One is the event horizon and we have chosen to put the origin of our spatial coordinate $y = 0$ at its location. There is also a smooth inner horizon, the Cauchy horizon, at some negative value of y .




FIG. 7: (a) Semiclassical static black hole solution. The numerical integration is started with initial conditions that ensure smoothness at the event horizon, which is located at the origin of the spatial coordinate. The fields Z , ξ and ψ all approach zero at a singularity inside the black hole. (b) The corresponding classical black hole solution with a smooth inner horizon. At the classical level there is no charge screening so the black hole carries fixed charge and there is no Z field.

The numerical solution of the semiclassical equations is started at $y = 0$ and the initial data chosen so as to obtain a black hole with a smooth event horizon [41]. To obtain a smooth inner horizon also requires some tuning of parameters which turns out, however, to be incompatible with the requirements for a smooth event horizon. In other words, we are not able to find static semiclassical solutions with two smooth horizons. Instead we encounter a singularity inside the black hole, where ψ , ξ and Z all go to zero. As we approach the singularity our numerical solution eventually breaks down but, as far as the numerics go, the indication is that we are dealing with a spacelike curvature singularity. The notion of a strong spacelike singularity is also supported by analytic scaling solutions, presented in [15], which we believe to be asymptotic to an exact solution sufficiently near the singularity.

The Cauchy horizon of a classical static black hole is replaced by a spacelike singularity at the semiclassical level. This can be traced to charge screening due to pair-production inside the black hole. For a static semiclassical solution the constraint equations (14) reduce to

$$\frac{d^2\psi}{dy^2} = -\frac{1}{2} \left(\frac{dZ}{dy} \right)^2. \quad (15)$$

As we go deeper into the black hole, *i.e.* towards negative y , any change in Z will focus

the area function towards zero faster than in the corresponding classical solution (where ψ is linear in y). As discussed above, charge screening appears as spatial variation in Z and therefore screening contributes to the focusing of the area function. This ultimately leads to the singular behavior seen in Figure 7a.

E. Neutral matter and the Hawking effect

So far we have considered semiclassical effects involving charged matter but we can also include electrically neutral matter along the lines of [32] by introducing conformally coupled scalar fields f_i , with flavor index $N = 1, \dots, N$,

$$S_f = -\frac{1}{2} \int d^2x \sqrt{-g} \sum_{i=1}^N (\nabla f_i)^2. \quad (16)$$

This gives rise to the following non-local term in the semiclassical effective action,

$$-\frac{N}{48} \int d^2x \sqrt{-g(x)} \int d^2x' \sqrt{-g(x')} R(x) G(x, x') R(x'), \quad (17)$$

in a leading $1/N$ approximation for $N \gg 1$, where $G(x, x')$ is a Green function for the ∇^2 operator. The analysis of semiclassical models of this type (see [42, 43, 44, 45, 46] for reviews) is usually carried out in conformal gauge where (17) reduces to

$$-2\kappa \int d^2x \partial_+ \rho \partial_- \rho, \quad (18)$$

with $\kappa \equiv N/12$. In [15] the so-called RST-term [47, 48],

$$-\frac{N}{48} \int d^2x \sqrt{-g} R \phi = -2\kappa \int d^2x \phi \partial_+ \partial_- \rho, \quad (19)$$

was also included in the action. In that case the model has exact semiclassical solutions for electrically neutral black holes which can be useful even if our main interest here is in charged black holes.

A closely related model was studied in [49], where analytic results were obtained for the combination of Hawking radiation and pair-production of charged particles in the background geometry of a classical two-dimensional charged black hole, but the semiclassical back-reaction on the geometry, which is the main concern of [15], was not considered.

The semiclassical equations become more complicated when the additional terms that come from (18) and (19) are included, but they can still be solved numerically. Both static

and dynamical black hole solutions were obtained in [15]. The qualitative nature of the solutions depends on the model parameters, in particular the relative size of κ and e^2 .

Let us first consider static black holes. For $e^2 \gg \kappa$ the static solutions are similar to the ones with $\kappa = 0$ that we have discussed already. As mentioned above, it is this limit that is of most interest if we want to model the physics of electrons with a large charge-to-mass ratio. In the opposite limit $\kappa \gg e^2$ the static solutions are quite different. There is still a curvature singularity where $\xi(y)$ goes to zero inside the black hole but the area function ψ reaches a local minimum before this singularity is reached. The area bounces back to large values and is divergent at the singularity. This means that the two-dimensional gravitational coupling goes to zero at the singularity and, from the 3+1-dimensional point of view, the area of the transverse two-sphere goes to infinity in a region of infinite spacetime curvature. This type of “bounce” singularity has been seen previously in static semiclassical solutions in two-dimensional dilaton gravity [29, 41, 50] but for physics the more interesting question is whether analogous behavior occurs in dynamical solutions where a black hole forms in gravitational collapse.

This brings us to dynamical solutions of the semiclassical equations with both charged and neutral matter effects included. This is a more challenging numerical problem than the one with only charged matter included, partly because the equations themselves are more complicated, but also due to subtleties involving boundary conditions that have to be imposed in the strong-coupling region [47, 48]. The numerical solutions reported in [15] show the formation and subsequent evaporation of a black hole with a spacelike singularity. In particular no bouncing of the area function is observed. It should be noted, however, that due to redshift effects a numerical code like the one employed in [15], which is based on a fixed coordinate grid, cannot resolve well the neighborhood of the event horizon at late stages in the evaporation of black hole that starts out large. This leaves open the possibility that the singularity inside a large charged black hole has a null portion close to the evaporation endpoint [51]. Presumably a more sophisticated numerical approach is needed to settle this issue.

Acknowledgments

We thank the organizers of the Workshop on Strings, Black Holes, and Quantum Space-time, Bokwang Phoenix Park, South Korea, April 21-23, 2006, for the opportunity to present this work, which was supported in part by grants from the Icelandic Research Fund and the University of Iceland Research Fund. Section III is based on work in collaboration with Andrei Frolov and Kristjan Kristjansson that appeared in [14, 15]. We would also like to thank Amos Ori for useful discussions.

-
- [1] R. Penrose, in in *Battelle Recontres* (ed. by C.M. DeWitt and J.A. Wheeler), W.A. Benjamin, New York, 121-235, (1968).
 - [2] M. Simpson and R. Penrose, *Int. J. Theor. Phys.* **7** 183 (1973).
 - [3] J.M. McNamara, *Proc. R. Soc. London* **A358** 499 (1978).
 - [4] I.D. Novikov and A.A. Starobinsky, *Sov. Phys. JETP* **51** 1 (1980).
 - [5] S. Chandrasekhar and J.B. Hartle, *Proc. R. Soc. London* **A384** 301 (1982).
 - [6] N. Zamorano, *Phys. Rev. D* **26** 2564 (1982).
 - [7] E. Poisson and W. Israel, *Phys. Rev. D* **41**, 1796 (1990).
 - [8] A. Ori, *Phys. Rev. Lett.* **67** 798 (1991).
 - [9] P. R. Brady and J. D. Smith, *Phys. Rev. Lett.* **75**, 1256 (1995) [arXiv:gr-qc/9506067].
 - [10] S. Hod and T. Piran, *Phys. Rev. Lett.* **81**, 1554 (1998) [arXiv:gr-qc/9803004].
 - [11] M. Dafermos, arXiv:gr-qc/0307013.
 - [12] R. Herman and W. A. Hiscock, *Phys. Rev. D* **49**, 3946 (1994).
 - [13] S. W. Hawking, *Phys. Rev. D* **14**, 2460 (1976).
 - [14] A. V. Frolov, K. R. Kristjansson and L. Thorlacius, *Phys. Rev. D* **72**, 021501 (2005) [arXiv:hep-th/0504073].
 - [15] A. V. Frolov, K. R. Kristjansson and L. Thorlacius, *Phys. Rev. D* **73**, 124036 (2006) [arXiv:hep-th/0604041].
 - [16] L. Susskind, L. Thorlacius and J. Uglum, *Phys. Rev. D* **48**, 3743 (1993) [arXiv:hep-th/9306069].
 - [17] D. N. Page, *Phys. Rev. Lett.* **44**, 301 (1980).

- [18] G. 't Hooft, Phys. Scripta **T36**, 247 (1991), and references therein.
- [19] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, Phys. Rev. D **55**, 5112 (1997) [arXiv:hep-th/9610043].
- [20] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [21] A. Strominger and C. Vafa, Phys. Lett. B **379**, 99 (1996) [arXiv:hep-th/9601029].
- [22] D. A. Lowe and L. Thorlacius, Phys. Rev. D **73**, 104027 (2006) [arXiv:hep-th/0601059].
- [23] S. B. Giddings, arXiv:hep-th/0605196.
- [24] W.K. Wothers and W.H. Zurek, Nature **299**, 802 (1982).
- [25] D. Dieks, Phys. Lett. A **92**, 271 (1982).
- [26] L. Susskind and L. Thorlacius, Phys. Rev. D **49**, 966 (1994) [arXiv:hep-th/9308100].
- [27] D. N. Page, Phys. Rev. Lett. **71**, 1291 (1993) [arXiv:gr-qc/9305007].
- [28] X. H. Ge and Y. G. Shen, Int. J. Mod. Phys. D **14**, 1321 (2005) [arXiv:gr-qc/0503094].
- [29] R. Balbinot and P. R. Brady, Class. Quant. Grav. **11**, 1763 (1994).
- [30] J. S. F. Chan and R. B. Mann, Phys. Rev. D **50**, 7376 (1994) [arXiv:gr-qc/9406021].
- [31] S. Droz, Phys. Lett. A **191**, 211 (1994).
- [32] C. G. . Callan, S. B. Giddings, J. A. Harvey and A. Strominger, Phys. Rev. D **45**, 1005 (1992) [arXiv:hep-th/9111056].
- [33] S. B. Giddings and A. Strominger, Phys. Rev. D **46**, 627 (1992) [arXiv:hep-th/9202004].
- [34] T. Banks, A. Dabholkar, M. R. Douglas and M. O'Loughlin, Phys. Rev. D **45**, 3607 (1992) [arXiv:hep-th/9201061].
- [35] J.S. Schwinger, Phys. Rev. **82**, 664 (1951).
- [36] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975) [Erratum-ibid. **46**, 206 (1976)].
- [37] S. R. Coleman, R. Jackiw and L. Susskind, Annals Phys. **93**, 267 (1975).
- [38] S. R. Coleman, Annals Phys. **101**, 239 (1976).
- [39] D. Grumiller and R. Meyer, arXiv:hep-th/0607030.
- [40] J. G. Russo, L. Susskind and L. Thorlacius, Phys. Lett. B **292**, 13 (1992) [arXiv:hep-th/9201074].
- [41] B. Birnir, S. B. Giddings, J. A. Harvey and A. Strominger, Phys. Rev. D **46**, 638 (1992) [arXiv:hep-th/9203042].
- [42] S. B. Giddings, arXiv:hep-th/9412138.

- [43] A. Strominger, arXiv:hep-th/9501071.
- [44] L. Thorlacius, Nucl. Phys. Proc. Suppl. **41**, 245 (1995) [arXiv:hep-th/9411020].
- [45] D. Grumiller, W. Kummer and D. V. Vassilevich, Phys. Rept. **369**, 327 (2002) [arXiv:hep-th/0204253].
- [46] A. Fabbri and J. Navarro-Salas, *Modeling black hole evaporation* (Imperial College Press, London, UK, 2005).
- [47] J. G. Russo, L. Susskind and L. Thorlacius, Phys. Rev. D **46**, 3444 (1992) [arXiv:hep-th/9206070].
- [48] J. G. Russo, L. Susskind and L. Thorlacius, Phys. Rev. D **47**, 533 (1993) [arXiv:hep-th/9209012].
- [49] A. Ori, Phys. Rev. D **63**, 104016 (2001) [arXiv:gr-qc/0102067].
- [50] L. Susskind and L. Thorlacius, Nucl. Phys. B **382**, 123 (1992) [arXiv:hep-th/9203054].
- [51] A. Ori, private communication (2006).